

Novel carboxylic acid derivatives, their preparation and use

The present invention relates to novel carboxylic acid derivatives, their preparation and use.


Endothelin is a peptide which is composed of 21 amino acids and is synthesized and released by the vascular endothelium. Endothelin exists in three isoforms, ET-1, ET-2 and ET-3. In the following text, "endothelin" or "ET" signifies one or all isoforms of endothelin. Endothelin is a potent vasoconstrictor and has a potent effect on vessel tone. It is known that this vasoconstriction is caused by binding of endothelin to its receptor (Nature, 332, (1988) 411-415; FEBS Letters, 231, (1988) 440-444 and Biochem. Biophys. Res. Commun., 154, (1988) 868-875).

Increased or abnormal release of endothelin causes persistent vasoconstriction in the peripheral, renal and cerebral blood vessels, which may lead to illnesses. It has been reported in the literature that elevated plasma levels of endothelin were found in patients with hypertension, acute myocardial infarct, pulmonary hypertension, Raynaud's syndrome, atherosclerosis and in the airways of asthmatics (Japan J. Hypertension, 12, (1989) 79, J. Vascular Med. Biology 2, (1990) 207, J. Am. Med. Association 264, (1990) 2868).

Accordingly, substances which specifically inhibit the binding of endothelin to the receptor ought also to antagonize the various abovementioned physiological effects of endothelin and therefore be valuable drugs.

We have found that certain carboxylic acid derivatives are good inhibitors of endothelin receptors.

The invention relates to carboxylic acid derivatives of the formula I



where R is formyl, tetrazole [sic], nitrile [sic], a COOH group or a radical which can be hydrolyzed to COOH, and the other substituents have the following meanings:

- 5 R² hydrogen, hydroxyl, NH₂, NH(C₁-C₄-alkyl), N(C₁-C₄-alkyl)₂, halogen, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy or C₁-C₄-alkylthio;
- 10 X nitrogen or CR¹⁴ where R¹⁴ is hydrogen or C₁-5-alkyl, or CR¹⁴ forms together with CR³ a 5- or 6-membered alkylene or alkenylene ring which can be substituted by one or two C₁-4-alkyl groups and in which in each case a methylene group can be replaced by oxygen, sulfur, -NH or -NC₁-4-alkyl;
- 15 R³ hydrogen, hydroxyl, NH₂, NH(C₁-C₄-Alkyl), N(C₁-C₄-alkyl)₂, halogen, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy, -NH-O-C₁-4-alkyl, C₁-C₄-alkylthio or CR³ is linked to CR¹⁴ as indicated above to give a 5- or 6-membered ring;
- 20 R⁴ and R⁵ (which can be identical or different):
- 25 phenyl or naphthyl, which can be substituted by one or more of the following radicals: halogen, nitro, cyano, hydroxyl, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy, phenoxy, C₁-C₄-alkylthio, amino, C₁-C₄-alkylamino or C₁-C₄-di-alkylamino; or
- 30 phenyl or naphthyl, which are connected together in the ortho positions via a direct linkage, a methylene, ethylene or ethenylene group, an oxygen or sulfur atom or an SO₂-, NH- or N-alkyl group, or C₃-C₇-cycloalkyl;
- 35 R⁶ hydrogen, C₁-C₈-alkyl, C₃-C₆-alkenyl, C₃-C₆-alkynyl or C₃-C₈-cycloalkyl, where each of these radicals can be substituted one or more times by: halogen, nitro, cyano, C₁-C₄-alkoxy, C₃-C₆-alkenyloxy, C₃-C₆-alkynyloxy, C₁-C₄-alkylthio, C₁-C₄-haloalkoxy, C₁-C₄-alkylcarbonyl, C₁-C₄-alkoxy-carbonyl, C₃-8-alkylcarbonylalkyl, C₁-C₄-alkylamino,
- 40 di-C₁-C₄-alkylamino, phenyl or phenyl or phenoxy which is substituted one or more times, eg. one to three times, by halogen, nitro, cyano, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy or C₁-C₄-alkylthio;
- 45 phenyl or naphthyl, each of which can be substituted by one or more of the following radicals: halogen, nitro, cyano, hydroxyl, amino, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy,

C₁-C₄-haloalkoxy, phenoxy, C₁-C₄-alkylthio, C₁-C₄-alkylamino, C₁-C₄-dialkylamino, dioxomethylene [sic] or dioxoethylene [sic];

5 a five- or six-membered heteroaromatic moiety containing one to three nitrogen atoms and/or one sulfur or oxygen atom, which can carry one to four halogen atoms and/or one or two of the following radicals: C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy, C₁-C₄-alkylthio, phenyl,
10 phenoxy or phenylcarbonyl, it being possible for the phenyl radicals in turn to carry one to five halogen atoms and/or one to three of the following radicals: C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy and/or C₁-C₄-alkylthio;

15 with the proviso that R⁶ can be hydrogen only when Z is not a single bond;

Y sulfur or oxygen or a single bond;

20

Z sulfur or oxygen or a single bond.


The compounds, and the intermediates for preparing them, such as IV and VI, may have one or more asymmetrical substituted carbon
25 atoms. Such compounds may be in the form of the pure enantiomers or pure diastereomers or a mixture thereof. The use of an enantiomerically pure compound as active substance is preferred.

The invention furthermore relates to the use of the abovementioned
30 carboxylic acid derivatives for producing drugs, in particular for producing endothelin receptor inhibitors.

The invention furthermore relates to the preparation of the compounds of the formula IV in enantiomerically pure form. Enantio-
35 selective epoxidation of an olefin with two phenyl substituents is known (J. Org. Chem. 59, 1994, 4378-4380). We have now found, surprisingly, that even ester groups in these systems permit epoxidation in high optical purity.


40 The preparation of the compounds according to the invention where Z is sulfur or oxygen starts from the epoxides IV, which are obtained in a conventional manner, eg. as described in J. March, Advanced Organic Chemistry, 2nd ed., 1983, page 862 and page 750, from the ketones II or the olefins III:

45



15 Carboxylic acid derivatives of the general formula VI can be prepared by reacting the epoxides of the general formula IV (eg. with $R = ROOR^{10}$ [sic]) with alcohols or thiols of the general formula V where R^6 and Z have the meanings stated in claim 1.

20



25

To do this, compounds of the general formula IV are heated with compounds of the formula V, in the molar ratio of about 1:1 to 1:7, preferably 1 to 3 mole equivalents, to 50-200°C, preferably 80-150°C.

30

The reaction can also take place in the presence of a diluent. All solvents which are inert toward the reagents used can be used for this purpose.

35 Examples of such solvents or diluents are water, aliphatic, alicyclic and aromatic hydrocarbons, which may in each case be chlorinated, such as hexane, cyclohexane, petroleum ether, naphtha, benzene, toluene, xylene, methylene chloride, chloroform, carbon tetrachloride, ethyl chloride and trichloroethylene, ethers such as diisopropyl ether, dibutyl ether, methyl tert-butyl ether, 40 propylene oxide, dioxane and tetrahydrofuran, ketones such as acetone, methyl ethyl ketone, methyl isopropyl ketone and methyl isobutyl ketone, nitriles such as acetonitrile and propionitrile, alcohols, such as methanol, ethanol, isopropanol, butanol and 45 ethylene glycol, esters such as ethyl acetate and amyl acetate, amides such as dimethylformamide, dimethylacetamide and N-methylpyrrolidone, sulfoxides and sulfones, such as dimethyl sulfoxide.

and sulfolane, bases such as pyridine, cyclic ureas such as 1,3-dimethylimidazolidin-2-one and 1,3-dimethyl-3,4,5,6-tetrahydro-2(1H)-pyrimidinone.

- 5 The reaction is preferably carried out at a temperature in the range from 0°C to the boiling point of the solvent or mixture of solvents.

The presence of a catalyst may be advantageous. Suitable catalysts are strong organic and inorganic acids, and Lewis acids. Examples thereof are, inter alia, sulfuric acid, hydrochloric acid, trifluoroacetic acid, p-toluenesulfonic acid, boron trifluoride etherate and titanium(IV) alcoholates.

- 15 Compounds of the formula VI where R⁴ and R⁵ are cycloalkyl can also be prepared by subjecting compounds of the formula VI where R⁴ and R⁵ are phenyl, naphthyl, or phenyl or naphthyl substituted as described above, to a nuclear hydrogenation.


- 20 Compounds of the formula VI can be obtained in enantiomerically pure form by starting from enantiomerically pure compounds of the formula IV and reacting them in the manner described with compounds of the formula V.

- 25 It is furthermore possible to obtain enantiomerically pure compounds of the formula VI by carrying out a classical racemate resolution on racemic or diastereomeric compounds of the formula VI using suitable enantiomerically pure bases such as brucine, strychnine, quinine, quinidine, chinchonidine [sic], chinchonine

- 30 [sic], yohimbine, morphine, dehydroabietylamine, ephedrine (-), (+), deoxyephedrine (+), (-), threo-2-amino-1-(p-nitrophenyl)-1,3-propanediol (+), (-), threo-2-(N,N-dimethylamino)-1-(p-nitrophenyl)-1,3-propanediol (+), (-) threo-2-amino-1-phenyl-1,3-propanediol (+), (-), α-methylbenzylamine (+), (-),
 35 α-(1-naphthyl)ethylamine (+), (-), α-(2-naphthyl)ethylamine (+), (-), aminomethylpinane, N,N-dimethyl-1-phenylethylamine, N-methyl-1-phenylethylamine, 4-nitrophenylethylamine, pseudoephedrine, norephedrine, norpseudoephedrine, amino acid derivatives, peptide derivatives.

40

- The compounds according to the invention where Y is oxygen, and the remaining substituents have the meanings stated under the general formula I, can be prepared, for example, by reacting the carboxylic acid derivatives of the general formula VI where the
 45 substituents have the stated meanings with compounds of the general formula VII



10

where R¹⁵ is halogen or R¹⁶-SO₂-, where R¹⁶ can be C₁-C₄-alkyl, C₁-C₄-haloalkyl or phenyl. The reaction preferably takes place in one of the abovementioned inert diluents with the addition of a suitable base, ie. of a base which deprotonates the intermediate

15 VI, in a temperature range from room temperature to the boiling point of the solvent.

Compounds of the formula VII are known, some of them can be bought, or they can be prepared in a generally known manner.

20


It is possible to use as base an alkali metal or alkaline earth metal hydride such as sodium hydride, potassium hydride or calcium hydride, a carbonate such as an alkali metal carbonate, eg. sodium or potassium carbonate, an alkali metal or alkaline earth

25 metal hydroxide such as sodium or potassium hydroxide, an organo-metallic compound such as butyllithium, or an alkali metal amide such as lithium diisopropylamide.

The compounds according to the invention where Y is sulfur, and

30 the remaining substituents have the meanings stated under the general formula I, can be prepared, for example, by reacting carboxylic acid derivatives of the general formula VIII, which can be obtained in a known manner from compounds of the general formula VI and in which the substituents have the abovementioned

35 meanings, with compounds of the general formula IX, where R², R³ and X have the meanings stated under general formula I.




The reaction preferably takes place in one of the abovementioned inert diluents with the addition of a suitable base, ie. a base which deprotonates the intermediate IX, in a temperature range from room temperature to the boiling point of the solvent.

5

It is possible to use as base, besides those mentioned above, organic bases such as triethylamine, pyridine, imidazole or diazabicycloundecane [sic].

10 Carboxylic acid derivatives of the formula VIa (Z in formula VI = direct linkage) can be prepared by reacting epoxides of the formula IV with cuprates of the formula XI:

15



20

XI

VIa

The cuprates can be prepared as described in Tetrahedron Letters 23, (1982) 3755.


25

Compounds of the formula I can also be prepared by starting from the corresponding carboxylic acids, ie. compounds of the formula I where R is COOH, and initially converting these in a conventional manner into an activated form, such as a halide, an anhydride or imidazolide, and then reacting the latter with an appropriate hydroxy compound HOR¹⁰. This reaction can be carried out in the usual solvents and often requires addition of a base, in which case those mentioned above are suitable. These two steps can also be simplified, for example, by allowing the carboxylic acid to act on the hydroxy compound in the presence of a dehydrating agent such as a carbodiimide.

In addition, it is also possible for compounds of the formula I to be prepared by starting from the salts of the corresponding carboxylic acids, ie. from compounds of the formula I where R is COR¹ and R¹ is OM, where M can be an alkali metal cation or the equivalent of an alkaline earth metal cation. These salts can be reacted with many compounds of the formula R¹-A where A is a conventional nucleofugic leaving group, for example halogen such as chlorine, bromine, iodine or aryl- or alkylsulfonyl which is unsubstituted or substituted by halogen, alkyl or haloalkyl, such as toluenesulfonyl and methylsulfonyl, or another equivalent

leaving group. Compounds of the formula R^1-A with a reactive substituent A are known or can be easily obtained with general expert knowledge. This reaction can be carried out in conventional solvents and advantageously takes place with the addition of a base, in which case those mentioned above are suitable.

The radical R in formula I may vary widely. For example, R is a group



where R^1 has the following meanings:

a) hydrogen;

b) succinylimidoxy [sic];

c) a five-membered heteroaromatic moiety linked by a nitrogen atom, such as pyrrolyl, pyrazolyl, imidazolyl and triazolyl, which may carry one or two halogen atoms, in particular fluorine and chlorine and/or one or two of the following radicals:

C_1-C_4 -alkyl such as methyl, ethyl, 1-propyl, 2-propyl, 2-methyl-2-propyl, 2-methyl-1-propyl, 1-butyl, 2-butyl;

C_1-C_4 -haloalkyl, in particular C_1-C_2 -haloalkyl such as fluoromethyl, difluoromethyl, trifluoromethyl, chlorodifluoromethyl, dichlorofluoromethyl, trichloromethyl, 1-fluoroethyl, 2-fluoroethyl, 2,2-difluoroethyl, 2,2,2-trifluoroethyl, 2-chloro-2,2-difluoroethyl, 2,2-dichloro-2-fluoroethyl, 2,2,2-trichloroethyl and pentafluoroethyl;

C_1-C_4 -haloalkoxy, in particular C_1-C_2 -haloalkoxy such as difluoromethoxy, trifluoromethoxy, chlorodifluoromethoxy, 1-fluoroethoxy, 2-fluoroethoxy, 2,2-difluoroethoxy, 1,1,2,2-tetrafluoroethoxy, 2,2,2-trifluoroethoxy, 2-chloro-1,1,2-trifluoroethoxy and pentafluoroethoxy, in particular trifluoromethoxy;


C_1-C_4 -alkoxy such as methoxy, ethoxy, propoxy, 1-methylethoxy, butoxy, 1-methylpropoxy, 2-methylpropoxy, 1,1-dimethylethoxy, in particular methoxy, ethoxy, 1-methylethoxy;

C₁-C₄-alkylthio such as methylthio, ethylthio, propylthio, 1-methylethylthio, butylthio, 1-methylpropylthio, 2-methylpropylthio, 1,1-dimethylethylthio, in particular methylthio and ethylthio;

5

d) R¹ furthermore a radical

10



where m is 0 or 1 and R⁷ and R⁸, which can be identical or different, have the following meanings:

15

hydrogen

C₁-C₈-alkyl, in particular C₁-C₄-alkyl as mentioned above;

20

C₃-C₆-alkenyl such as 2-propenyl, 2-butenyl, 3-butenyl, 1-methyl-2-propenyl, 2-methyl-2-propenyl, 2-pentenyl, 3-pentenyl, 4-pentenyl, 1-methyl-2-butenyl, 2-methyl-2-butenyl, 3-methyl-2-butenyl, 1-methyl-3-butenyl,

25

2-methyl-3-butenyl, 3-methyl-3-butenyl, 1,1-dimethyl-2-propenyl, 1,2-dimethyl-2-propenyl, 1-ethyl-2-propenyl, 2-hexenyl, 3-hexenyl, 4-hexenyl, 5-hexenyl, 1-methyl-2-pentenyl, 2-methyl-2-pentenyl, 3-methyl-2-pentenyl, 4-methyl-2-pentenyl, 3-methyl-3-pentenyl, 4-methyl-3-pentenyl, 1-methyl-4-pentenyl, 2-methyl-4-pentenyl, 3-methyl-4-pentenyl, 4-methyl-4-pentenyl, 1,1-dimethyl-2-butenyl, 1,1-dimethyl-3-butenyl, 1,2-dimethyl-2-butenyl, 1,2-dimethyl-3-butenyl, 1,3-dimethyl-2-butenyl, 1,3-dimethyl-3-butenyl, 2,2-dimethyl-3-butenyl, 2,3-dimethyl-2-butenyl, 2,3-dimethyl-3-butenyl, 1-ethyl-2-butenyl, 1-ethyl-3-butenyl, 2-ethyl-2-butenyl, 2-ethyl-3-butenyl, 1,1,2-trimethyl-2-propenyl, 1-ethyl-1-methyl-2-propenyl and 1-ethyl-2-methyl-2-propenyl, in particular 2-propenyl, 2-butenyl, 3-methyl-2-butenyl and 3-methyl-2-pentenyl;

30

35

40

C₃-C₆-alkynyl such as 2-propynyl, 2-butyne, 3-butyne, 1-methyl-2-propynyl, 2-pentyne, 3-pentyne, 4-pentyne, 1-methyl-3-butyne, 2-methyl-3-butyne, 1-methyl-2-butyne, 1,1-dimethyl-2-propynyl, 1-ethyl-2-propynyl, 2-hexynyl, 3-hexynyl, 4-hexynyl, 5-hexynyl, 1-methyl-2-pentyne, 1-methyl-2-pentyne, 1-methyl-3-pentyne, 1-methyl-4-pentyne, 2-methyl-3-pentyne, 2-methyl-4-pentyne, 3-methyl-4-pentyne,

45

10

4-methyl-2-pentynyl, 1,1-dimethyl-2-butynyl,
 1,1-dimethyl-3-butynyl, 1,2-dimethyl-3-butynyl,
 2,2-dimethyl-3-butynyl, 1-ethyl-2-butynyl, 1-ethyl-3-butynyl,
 2-ethyl-3-butynyl and 1-ethyl-1-methyl-2-propynyl, preferably
 2-propynyl, 2-butynyl, 1-methyl-2-propynyl and
 1-methyl-2-butynyl, in particular 2-propynyl

C₃-C₈-cycloalkyl such as cyclopropyl, cyclobutyl, cyclopentyl,
 cyclohexyl and cycloheptyl, cyclooctyl, where these alkyl,
 cycloalkyl, alkenyl and alkynyl groups can each carry one to
 five halogen atoms, in particular fluorine or chlorine and/or
 one or two of the following groups:

C₁-C₄-alkyl, C₁-C₄-alkoxy, C₁-C₄-alkylthio, C₁-C₄-haloalkoxy as
 mentioned above, C₃-C₆-alkenyloxy, C₃-C₆-alkenylthio,
 C₃-C₆-alkynyloxy, C₃-C₆-alkynylthio, where the alkenyl and
 alkynyl constituents present in these radicals preferably
 have the abovementioned meanings;

C₁-C₄-alkylcarbonyl such as, in particular, methylcarbonyl,
 ethylcarbonyl, propylcarbonyl, 1-methylethylcarbonyl, butyl-
 carbonyl, 1-methylpropylcarbonyl, 2-methylpropylcarbonyl,
 1,1-dimethylethylcarbonyl;

C₁-C₄-alkoxycarbonyl such as methoxycarbonyl, ethoxycarbonyl,
 propyloxy carbonyl, 1-methylethoxycarbonyl, butyloxy carbonyl,
 1-methylpropyloxy carbonyl, 2-methylpropyloxy carbonyl,
 1,1-dimethylethoxycarbonyl;

C₃-C₆-alkenylcarbonyl, C₃-C₆-alkynylcarbonyl, C₃-C₆-alkenyloxy-
 carbonyl and C₃-C₆-alkynyloxy carbonyl, where the alkenyl and
 alkynyl radicals are preferably defined as detailed above;

phenyl, unsubstituted or substituted one or more times, eg.
 one to three times, by halogen, nitro, cyano, C₁-C₄-alkyl,
 C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy or C₁-C₄-alkyl-
 thio, such as 2-fluorophenyl, 3-chlorophenyl, 4-bromophenyl,
 2-methylphenyl, 3-nitrophenyl, 4-cyanophenyl, 2-trifluoro-
 methylphenyl, 3-methoxyphenyl, 4-trifluoroethoxyphenyl,
 2-methylthiophenyl, 2,4-dichlorophenyl, 2-methoxy-3-methyl-
 phenyl, 2,4-dimethoxyphenyl, 2-nitro-5-cyanophenyl,
 2,6-difluorophenyl;

di-C₁-C₄-alkylamino such as, in particular, dimethylamino, dipropylamino, N-propyl-N-methylamino, N-propyl-N-ethylamino, diisopropylamino, N-isopropyl-N-methylamino, N-isopropyl-N-ethylamino, N-isopropyl-N-propylamino;

5


R⁷ and R⁸ furthermore phenyl which can be substituted by one or more, eg. one to three, of the following radicals: halogen, nitro, cyano, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy or C₁-C₄-alkylthio, as mentioned above in particular;

10

or R⁷ and R⁸ together form a C₄-C₇-alkylene chain which is closed to form a ring, is unsubstituted or substituted, eg. substituted by C₁-C₄-alkyl, and may contain a heteroatom selected from the group consisting of oxygen, sulfur or nitrogen, such as -(CH₂)₄-, -(CH₂)₅-, -(CH₂)₆-, -(CH₂)₇-, -(CH₂)₂-O-(CH₂)₂-, -CH₂-S-(CH₂)₃-, -(CH₂)₂-O-(CH₂)₃-, -NH-(CH₂)₃-, -CH₂-NH-(CH₂)₂-, -CH₂-CH=CH-CH₂-, -CH=CH-(CH₂)₃-;

15

20 e) R¹ furthermore a group



25

where k is 0, 1 and 2, p is 1, 2, 3 and 4 and R⁹ is

C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₃-C₆-alkenyl, C₃-C₆-alkynyl or unsubstituted or substituted phenyl, as mentioned above in particular.

30

f) R¹ furthermore a radical OR¹⁰, where R¹⁰ is:

35

hydrogen, the cation of an alkali metal such as lithium, sodium, potassium or the cation of an alkaline earth metal such as calcium, magnesium and barium or an environmentally compatible organic ammonium ion such as tertiary C₁-C₄-alkyl-ammonium or the ammonium ion;

40

C₃-C₈-cycloalkyl as mentioned above, which may carry one to three C₁-C₄-alkyl groups;

45

C₁-C₈-alkyl such as, in particular, methyl, ethyl, propyl, 1-methylethyl, butyl, 1-methylpropyl, 2-methylpropyl, 1,1-dimethylethyl, pentyl, 1-methylbutyl, 2-methylbutyl, 3-methylbutyl, 1,2-dimethylpropyl, 1,1-dimethylpropyl, 2,2-dimethylpropyl, 1-ethylpropyl, hexyl, 1-methylpentyl,

2-methylpentyl, 3-methylpentyl, 4-methylpentyl, 1,2-dimethylbutyl, 1,3-dimethylbutyl, 2,3-dimethylbutyl, 1,1-dimethylbutyl, 2,2-dimethylbutyl, 3,3-dimethylbutyl, 1,1,2-trimethylpropyl, 1,2,2-trimethylpropyl, 1-ethylbutyl, 2-ethylbutyl, 1-ethyl-2-methylpropyl, which can carry one to five halogen atoms, in particular fluorine and chlorine and/or one of the following radicals:

C_1 - C_4 -alkoxy, C_1 - C_4 -alkylthio, cyano, C_1 - C_4 -alkylcarbonyl, C_3 - C_8 -cycloalkyl, C_1 - C_4 -alkoxycarbonyl, phenyl, phenoxy or phenylcarbonyl, where the aromatic radicals in turn can carry in each case one to five halogen atoms and/or one to three of the following radicals: nitro, cyano, C_1 - C_4 -alkyl, C_1 - C_4 -haloalkyl, C_1 - C_4 -alkoxy, C_1 - C_4 -haloalkoxy and/or C_1 - C_4 -alkylthio, as mentioned above in particular;

a C_1 - C_8 -alkyl as mentioned above, which can carry one to five halogen atoms, in particular fluorine and/or chlorine, and carries one of the following radicals: a 5-membered heteroaromatic moiety containing one to three nitrogen atoms, or a 5-membered heteroaromatic moiety containing a nitrogen atom and an oxygen or sulfur atom, which can carry one to four halogen atoms and/or one or two of the following radicals:

nitro, cyano, C_1 - C_4 -alkyl, C_1 - C_4 -haloalkyl, C_1 - C_4 -alkoxy, phenyl, C_1 - C_4 -haloalkoxy and/or C_1 - C_4 -alkylthio. Particular mention may be made of: 1-pyrazolyl, 3-methyl-1-pyrazolyl, 4-methyl-1-pyrazolyl, 3,5-dimethyl-1-pyrazolyl, 3-phenyl-1-pyrazolyl, 4-phenyl-1-pyrazolyl, 4-chloro-1-pyrazolyl, 4-bromo-1-pyrazolyl, 1-imidazolyl, 1-benzimidazolyl, 1,2,4-triazol-1-yl, 3-methyl-1,2,4-triazol-1-yl, 5-methyl-1,2,4-triazol-1-yl, 1-benzotriazolyl, 3-isopropyl-5-isoxazolyl, 3-methyl-5-isoxazolyl, 2-oxazolyl, 2-thiazolyl, 2-imidazolyl, 3-ethyl-5-isoxazolyl, 3-phenyl-5-isoxazolyl, 3-tert-butyl-5-isoxazolyl;

a C_2 - C_6 -alkyl group which carries one of the following radicals in position 2: C_1 - C_4 -alkoxyimino, C_3 - C_6 -alkynyloxyimino, C_3 - C_6 -haloalkenyloxyimino or benzyloxyimino;


a C_3 - C_6 -alkenyl or C_3 - C_6 -alkynyl group, it being possible for these groups in turn to carry one to five halogen atoms;

R^{10} furthermore a phenyl radical which can carry one to five halogen atoms and/or one to three of the following radicals: nitro, cyano, C_1 - C_4 -alkyl, C_1 - C_4 -haloalkyl, C_1 - C_4 -alkoxy,

C₁-C₄-haloalkoxy and/or C₁-C₄-alkylthio, as mentioned above in particular;

5 a 5-membered heteroaromatic moiety which is linked via a nitrogen atom, contains one to three nitrogen atoms and can carry one or two halogen atoms and/or one or two of the following radicals: C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, phenyl, C₁-C₄-haloalkoxy and/or C₁-C₄-alkylthio. Particular
10 mention may be made of: 1-pyrazolyl, 3-methyl-1-pyrazolyl, 4-methyl-1-pyrazolyl, 3,5-dimethyl-1-pyrazolyl, 3-phenyl-1-pyrazolyl, 4-phenyl-1-pyrazolyl, 4-chloro-1-pyrazolyl, 4-bromo-1-pyrazolyl, 1-imidazolyl, 1-benzimidazolyl, 1,2,4-triazol-1-yl, 3-methyl-1,2,4-triazol-1-yl, 5-methyl-1,2,4-triazol-1-yl, 1-benzotriazolyl,
15 3,4-dichloro-1-imidazolyl;

R¹⁰ furthermore a group




25 where R¹¹ and R¹², which can be identical or different, are:

30 C₁-C₈-alkyl, C₃-C₆-alkenyl, C₃-C₆-alkynyl, C₃-C₈-cycloalkyl, it being possible for these radicals to carry a C₁-C₄-alkoxy, C₁-C₄-alkylthio and/or an unsubstituted or substituted phenyl radical, as mentioned above in particular;

35 phenyl which can be substituted by one or more, eg. one to three, of the following radicals: halogen, nitro, cyano, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy or C₁-C₄-alkylthio, where these radicals are, in particular, those mentioned above;

40 or R¹¹ and R¹² together form a C₃-C₁₂-alkylene chain which can carry one to three C₁-C₄-alkyl groups and contain a heteroatom from the group consisting of oxygen, sulfur and nitrogen, as mentioned in particular for R⁷ and R⁸.

g) R¹ furthermore a radical



5


where R^{13} is:

10 C_1 - C_4 -alkyl, C_3 - C_6 -alkenyl, C_3 - C_6 -alkynyl, C_3 - C_8 -cycloalkyl as mentioned above in particular, it being possible for these radicals to carry a C_1 - C_4 -alkoxy, C_1 - C_4 -alkylthio and/or a phenyl radical as mentioned above;

15 phenyl, unsubstituted or substituted, in particular as mentioned above.

h) R^1 a radical

20



25

where R^{13} has the abovementioned meaning.

R can furthermore be:

30 tetrazole [sic] or nitrile [sic].

In respect of the biological effect, preferred carboxylic acid derivatives of the general formula I, both as pure enantiomers and pure diastereomers or as mixture thereof, are those where the
35 substituents have the following meanings:

R^2 hydrogen, hydroxyl, $N(C_1$ - C_4 -alkyl) $_2$, the C_1 - C_4 -alkyl, C_1 - C_4 -haloalkyl, C_1 - C_4 -alkoxy, C_1 - C_4 -haloalkoxy, C_1 - C_4 -alkylthio groups and halogen atoms mentioned in detail
40 for R^1 , especially chlorine, methyl, methoxy, ethoxy, difluoromethoxy, trifluoromethoxy;

X nitrogen or CR^{14} where

45 R^{14} is hydrogen or alkyl, or CR^{14} forms together with CR^3 a 4- to 5-membered alkylene or alkenylene ring in which, in each case, a methylene group can be replaced by oxygen or sulfur,

such as $-\text{CH}_2-\text{CH}_2-\text{O}-$, $-\text{CH}=\text{CH}-\text{O}-$, $-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{O}-$, $-\text{CH}=\text{CH}-\text{CH}_2\text{O}-$,
 in particular hydrogen, $-\text{CH}_2-\text{CH}_2-\text{O}-$, $-\text{CH}(\text{CH}_3)-\text{CH}(\text{CH}_3)-\text{O}-$,
 $-\text{C}(\text{CH}_3)=\text{C}(\text{CH}_3)-\text{O}-$, $-\text{CH}=\text{C}(\text{CH}_3)-\text{O}-$ or $-\text{C}(\text{CH}_3)=\text{C}(\text{CH}_3)-\text{S}$;

5 R^3 the hydrogen, hydroxyl, $\text{N}(\text{C}_1-\text{C}_4\text{-alkyl})_2$, $\text{C}_1-\text{C}_4\text{-alkyl}$,
 $\text{C}_1-\text{C}_4\text{-haloalkyl}$, $\text{C}_1-\text{C}_4\text{-alkoxy}$, $\text{C}_1-\text{C}_4\text{-haloalkoxy}$, $\text{C}_1-\text{C}_4\text{-alkyl-}$
 thio groups and halogen atoms mentioned for R^1 , especially
 chlorine, methyl, methoxy, ethoxy, difluoromethoxy, tri-
 fluoromethoxy or is linked to R^{14} as mentioned above to give a
 10 5- or 6-membered ring;

R^4 and R^5 phenyl or naphthyl, which can be substituted by one or
 more, eg. one to three, of the following radicals: halogen,
 nitro, cyano, hydroxyl, mercapto, amino, $\text{C}_1-\text{C}_4\text{-alkyl}$,
 15 $\text{C}_1-\text{C}_4\text{-haloalkyl}$, $\text{C}_1-\text{C}_4\text{-alkoxy}$, $\text{C}_1-\text{C}_4\text{-haloalkoxy}$,
 $\text{C}_1-\text{C}_4\text{-alkylthio}$, $\text{C}_1-\text{C}_4\text{-alkylamino}$, di- $\text{C}_1-\text{C}_4\text{-alkylamino}$,
 $\text{C}_1-\text{C}_4\text{-alkylcarbonyl}$, $\text{C}_1-\text{C}_4\text{-alkoxycarbonyl}$; phenyl or naphthyl,
 which are connected together in the ortho positions by a
 direct linkage, a methylene, ethylene or ethenylene group,
 20 an oxygen or sulfur atom or an SO_2 , NH or N-alkyl group, or
 $\text{C}_3-\text{C}_7\text{-cycloalkyl}$;

R^6 $\text{C}_1-\text{C}_8\text{-alkyl}$, $\text{C}_3-\text{C}_6\text{-alkenyl}$, $\text{C}_3-\text{C}_6\text{-alkynyl}$ or $\text{C}_3-\text{C}_8\text{-cycloalkyl}$
 as mentioned above in particular, it being possible for these
 25 radicals in each case to be substituted one or more times by:
 halogen, hydroxyl, nitro, cyano, $\text{C}_1-\text{C}_4\text{-alkoxy}$, $\text{C}_3-\text{C}_6\text{-alkenyl-}$
 oxy , $\text{C}_3-\text{C}_6\text{-alkynyloxy}$, $\text{C}_1-\text{C}_4\text{-alkylthio}$, $\text{C}_1-\text{C}_4\text{-haloalkoxy}$,
 $\text{C}_1-\text{C}_4\text{-alkylcarbonyl}$, hydroxycarbonyl, $\text{C}_1-\text{C}_4\text{-alkoxycarbonyl}$,
 $\text{C}_1-\text{C}_4\text{-alkylamino}$, di- $\text{C}_1-\text{C}_4\text{-alkylamino}$ or unsubstituted or
 30 substituted phenyl or phenoxy, as mentioned above in
 particular;

phenyl or naphthyl, which can be substituted by one or more
 of the following radicals: halogen, nitro, cyano, hydroxyl,
 35 amino, $\text{C}_1-\text{C}_4\text{-alkyl}$, $\text{C}_1-\text{C}_4\text{-haloalkyl}$, $\text{C}_1-\text{C}_4\text{-alkoxy}$, $\text{C}_1-\text{C}_4\text{-halo-}$
 alkoxy , phenoxy, $\text{C}_1-\text{C}_4\text{-alkylthio}$, $\text{C}_1-\text{C}_4\text{-alkylamino}$ [sic] or
 $\text{C}_1-\text{C}_4\text{-dialkylamino}$, as mentioned in particular for R^7 and R^4 ;

a five- or six-membered heteroaromatic moiety which contains
 40 one to three nitrogen atoms and/or one sulfur or oxygen atom
 and which can carry one to four halogen atoms and/or one or
 two of the following radicals: $\text{C}_1-\text{C}_4\text{-alkyl}$, $\text{C}_1-\text{C}_4\text{-haloalkyl}$,
 $\text{C}_1-\text{C}_4\text{-alkoxy}$, $\text{C}_1-\text{C}_4\text{-haloalkoxy}$, $\text{C}_1-\text{C}_4\text{-alkylthio}$, phenyl,
 phenoxy or phenylcarbonyl, it being possible for the phenyl
 45 radicals in turn to carry one to five halogen atoms and/or
 one to three of the following radicals: $\text{C}_1-\text{C}_4\text{-alkyl}$,

C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy and/or C₁-C₄-alkylthio, as mentioned for R⁴ in particular;

Y sulfur, oxygen or a single bond;

5

Z sulfur, oxygen, -SO-, -SO₂- or a single bond.

Particularly preferred compounds of the formula I, both as pure enantiomers and pure diastereomers or as mixture thereof, are those in which the substituents have the following meanings:

10

R² C₁-C₄-alkyl, C₁-C₄-alkoxy

X nitrogen or CR¹⁴, where

15

R¹⁴ is hydrogen or alkyl, or CR¹⁴ forms together with CR³ a 4- or 5-membered alkylene or alkenylene ring such as -CH₂-CH₂-CH₂-, -CH=CH-CH₂-, in which in each case a methylene group can be replaced by oxygen or sulfur, such as -CH₂-CH₂-O-, -CH=CH-O-, -CH₂-CH₂-CH₂-O-, -CH=CH-CH₂O-, in particular hydrogen, -CH₂-CH₂-O-, -CH(CH₃)-CH(CH₃)-O-, -C(CH₃)=C(CH₃)-O-, -CH=C(CH₃)-O- or -C(CH₃)=C(CH₃)-S;

20

R³ the C₁-C₄-alkyl, C₁-C₄-alkoxy, C₁-C₄-alkylthio groups mentioned for R¹, or is linked to R¹⁴ as mentioned above to give a 5- or 6-membered ring;

25

R⁴ and R⁵ phenyl (identical or different) which can be substituted by one or more, eg. one to three, of the following radicals: halogen, nitro, hydroxyl, C₁-C₄-alkyl, C₁-C₄-alkoxy, C₁-C₄-alkylthio or

30

R⁴ and R⁵ are phenyl groups which are connected together in the ortho positions by a direct linkage, a methylene, ethylene or ethenylene group, an oxygen or sulfur atom or an SO₂, NH or N-alkyl group; or

35

R⁴ and R⁵ are C₃-C₇-cycloalkyl;

40

R⁶ C₁-C₈-alkyl, C₃-C₆-alkenyl or C₃-C₈-cycloalkyl, it being possible for these radicals in each case to be substituted one or more times by: halogen, hydroxyl, nitro, cyano, C₁-C₄-alkoxy, C₃-C₆-alkenyloxy, C₁-C₄-alkylthio;

45

phenyl or naphthyl, which can be substituted by one or more of the following radicals: halogen, nitro, cyano, hydroxyl, amino, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy, phenoxy, C₁-C₄-alkylthio, C₁-C₄-alkylamino [sic] or C₁-C₄-dialkylamino;

a five- or six-membered heteroaromatic moiety which contains a nitrogen atom and/or a sulfur or oxygen atom and which can carry one to four halogen atoms and/or one or two of the following radicals: C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-alkylthio, phenyl, phenoxy or phenylcarbonyl, it being possible for the phenyl radicals in turn to carry one to five halogen atoms and/or one to three of the following radicals: C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy and/or C₁-C₄-alkylthio;

Y sulfur, oxygen or a single bond;

Z sulfur, oxygen, -SO-, -SO₂- or a single bond.

The compounds of the present invention provide a novel therapeutic potential for the treatment of hypertension, pulmonary hypertension, myocardial infarct, angina pectoris, acute kidney failure, renal insufficiency, cerebral vasospasms, cerebral ischemia, subarachnoid hemorrhages, migraine, asthma, atherosclerosis, endotoxic shock, endotoxin-induced organ failure, intravascular coagulation, restenosis after angioplasty, benign prostatic hyperplasia, or hypertension or kidney failure caused by ischemia or intoxication.

The good effect of the compounds can be shown in the following tests:

Receptor binding studies

Cloned human ET_A receptor-expressing CHO cells and guinea pig cerebellar membranes with > 60 % ET_B compared with ET_A receptors were used for binding studies.

The ET_A receptor-expressing CHO cells were grown in F₁₂ medium containing 10 % fetal calf serum, 1 % glutamine, 100 U/ml penicillin and 0.2 % streptomycin (Gibco BRL, Gaithersburg, MD, USA). After 48 h, the cells were washed with PBS and incubated with 0.05 % trypsin-containing PBS for 5 min. Neutralization was then carried out with F₁₂ medium, and the cells were collected by centrifugation at 300 x g. To lyse the cells, the pellet was briefly washed with lysis buffer (5 mM Tris-HCl, pH 7.4 with 10 % gly-

cerol) and then incubated at a concentration of 10^7 cells/ml of lysis buffer at 4°C for 30 min. The membranes were centrifuged at $20,000 \times g$ for 10 min, and the pellet was stored in liquid nitrogen.

5

Guinea pig cerebella were homogenized in a Potter-Elvehjem homogenizer and [lacuna] obtained by differential centrifugation at $1000 \times g$ for 10 min and repeated centrifugation of the supernatant at $20,000 \times g$ for 10 min.

10

Binding assays

For the ET_A and ET_B receptor binding assay, the membranes were suspended in incubation buffer (50 mM Tris-HCl, pH 7.4 with 5 mM MnCl_2 , 40 $\mu\text{g/ml}$ bacitracin and 0.2 % BSA) at a concentration of 50 μg of protein per assay mixture and incubated with 25 pM [^{125}I]- ET_1 (ET_A receptor assay) or 25 pM [^{125}I]- RZ_3 (ET_B receptor assay) in the presence and absence of test substance at 25°C . The nonspecific binding was determined using 10^{-7} M ET_1 . After 30 min, the free and bound radioligand were separated by filtration through GF/B glass fiber filters (Whatman, England) on a Skatron cell collector (Skatron, Lier, Norway) and the filters were washed with ice-cold Tris-HCl buffer, pH 7.4 with 0.2 % BSA. The radioactivity collected on the filters was quantified using a Packard 2200 CA liquid scintillation counter.

Functional in vitro assay system to look for endothelin receptor (subtype A) antagonists

This assay system is a functional, cell-based assay for endothelin receptors. When certain cells are stimulated with endothelin 1 (ET_1) they show an increase in the intracellular calcium concentration. This increase can be measured in intact cells loaded with calcium-sensitive dyes.

35

1-Fibroblasts which had been isolated from rats and in which an endogenous endothelin receptor of the A subtype had been detected were loaded with the fluorescent dye Fura 2-am as follows: after trypsinization, the cells were resuspended in buffer A (120 mM NaCl, 5 mM KCl, 1.5 mM MgCl_2 , 1 mM CaCl_2 , 25 mM HEPES, 10 mM glucose, pH 7.4) to a density of $2 \times 10^6/\text{ml}$ and incubated with Fura 2-am (2 μM), Pluronic F-127 (0.04 %) and DMSO (0.2 %) at 37°C in the dark for 30 min. The cells were then washed twice with buffer A and resuspended at $2 \times 10^6/\text{ml}$.

45

The fluorescence signal from 2×10^5 cells per ml with Ex/Em 380/510 was recorded continuously at 30°C. The test substances and, after an incubation time of 3 min, ET1 [lacuna] to the cells, the maximum change in the fluorescence was determined. The response of the cells to ET1 without previous addition of a test substance was used as control and was set equal to 100 %.

Testing of ET antagonists in vivo

10 Male SD rats weighting 250-300 g were anesthetized with amobarbital, artificially ventilated, vagotomized and pithed. The carotid artery and jugular vein were cathetized [sic].

In control animals, intravenous administration of 1 µg/kg ET1 led to a distinct rise in blood pressure which persisted for a lengthy period.

The test animals received an i.v. injection of the test compounds (1 ml/kg) 5 min before the administration of ET1. To determine the ET-antagonistic properties, the rise in blood pressure in the test animals was compared with that in the control animals.

Endothelin-1-induced sudden death in mice

25 The principle of the test is the inhibition of the sudden heart death caused in mice by endothelin, which is probably induced by constriction of the coronary vessels, by pretreatment with endothelin receptor antagonists. Intravenous injection of 10 nmol/kg endothelin in a volume of 5 ml/kg of body weight results in death of the animals within a few minutes.

The lethal endothelin-1 dose is checked in each case on a small group of animals. If the test substance is administered intravenously, the endothelin-1 injection which was lethal in the reference group usually takes place 5 min thereafter. With other modes of administration, the times before administration are extended, where appropriate up to several hours.

The survival rate is recorded, and effective doses which protect 50 % of the animals (ED 50) from endothelin-induced heart death for 24 h or longer are determined.

Functional test on vessels for endothelin receptor antagonists

Segments of rabbit aorta are, after an initial tension of 2 g and a relaxation time of 1 h in Krebs-Henseleit solution at 37°C and pH 7.3-7.4, first induced to contract with K⁺. After washing out, an endothelin dose-effect plot up to the maximum is constructed.

5

Potential endothelin antagonists are administered to other preparations of the same vessel 15 min before starting the endothelin dose-effect plot. The effects of the endothelin are calibrated as a % of the K⁺-induced contraction. Effective endothelin antagonists result in a shift to the right in the endothelin dose-effect plot.

The compounds according to the invention can be administered orally or parenterally (subcutaneously, intravenously, intramuscularly, intraperitoneally) in a conventional way. Administration can also take place with vapors or sprays through the nasopharyngeal space.

The dosage depends on the age, condition and weight of the patient and on the mode of administration. The daily dose of active substance is, as a rule, about 0.5-50 mg/kg of body weight on oral administration and about 0.1-10 mg/kg of body weight on parenteral administration.

The novel compounds can be used in conventional solid or liquid pharmaceutical forms, eg. as uncoated or (film-)coated tablets, capsules, powders, granules, suppositories, solutions, ointments, creams or sprays. These are produced in a conventional way. The active substances can for this purpose be processed with conventional pharmaceutical aids such as tablet binders, fillers, preservatives, tablet disintegrants, flow regulators, plasticizers, wetting agents, dispersants, emulsifiers, solvents, release-slowing agents, antioxidants and/or propellant gases (cf. H. Sucker et al.: Pharmazeutische Technologie, Thieme-Verlag, Stuttgart, 1991). The administration forms obtained in this way normally contain from 0.1 to 90 % by weight of the active substance.

Synthesis examples

40

Example 1

Methyl 2-hydroxy-3-methoxy-3,3-diphenylpropionate

5 g (19.6 mmol) of methyl 3,3-diphenyl-2,3-epoxypropionate were dissolved in 50 ml of absolute methanol and, at 0°C, 0.1 ml of boron trifluoride etherate was added. The mixture was stirred at 0°C for 2 h and at room temperature for a further 12 h. The sol-

vent was distilled out, the residue was taken up in ethyl acetate, washed with sodium bicarbonate solution and water and dried over magnesium sulfate. After removal of the solvent by distillation there remained 5.5 g (88 %) of a pale yellow oil.

5

Example 2

Methyl 2-hydroxy-3-phenoxy-3,3-diphenylpropionate

5 g (19.6 mmol) of methyl 3,3-diphenyl-2,3-epoxypropionate and
10 5.6 g (60 mmol) of phenol were heated together at 100°C for 6 h. Removal of the excess phenol by distillation under high vacuum and purification of the residue by chromatography on silica gel with hexane/ethyl acetate mixtures resulted in 4.9 g (77 %) of a pale yellow oil.

15

Example 3

Methyl 2-(4,6-dimethoxy-pyrimidin-2-yloxy)-3-methoxy-3,3-diphenylpropionate

20 2.86 g (10 mmol) of methyl 2-hydroxy-3-methoxy-3,3-diphenylpropionate were dissolved in 40 ml of dimethylformamide, and 0.3 g (12 mmol) of sodium hydride was added. The mixture was stirred for 1 h and then 2.2 g (10 mmol) of 4,6-dimethoxy-2-methylsulfonylpyrimidine were added. After stirring at room temperature for 24 h, cautious hydrolysis was carried out with 10 ml of
25 water, the pH was adjusted to 5 with acetic acid, and the solvent was removed by distillation under high vacuum. The residue was taken up in 100 ml of ethyl acetate, washed with water and dried over magnesium sulfate, and the solvent was distilled out. The
30 residue was mixed with 10 ml of ether, and the resulting precipitate was filtered off with suction. After drying, 3.48 g (82 %) of a white powder remained.
Melting point 81°C.

35 Example 4

2-(4,6-Dimethoxy-pyrimidin-2-yloxy)-3-methoxy-3,3-diphenylpropionic acid

2.12 g (5 mmol) of methyl 2-(4,6-dimethoxy-pyrimidin-2-yl-
40 oxy)-3-methoxy-3,3-diphenylpropionate were dissolved in 50 ml of dioxane, 10 ml of 1 N KOH solution were added, and the mixture was stirred at 100°C for 3 h. The solution was diluted with 300 ml of water and extracted with ethyl acetate to remove unreacted ester. The aqueous phase was then adjusted to pH 1-2 with dilute
45 hydrochloric acid and extracted with ethyl acetate. After drying over magnesium sulfate and removal of the solvent by distillation, the residue was mixed with an ether/hexane mixture, and the

precipitate which formed was filtered off with suction. After drying, 1.85 g (90 %) of a white powder remained.

Melting point 167°C

5 Example 5

2-(4,6-Dimethoxy-2-pyrimidinyloxy)-3-methoxy-3,3-diphenyl sodium [sic] propionate

1.68 g (4 mmol) of 2-(4,6-dimethoxy-2-pyrimidinyloxy)-3-methoxy-3,3-diphenylpropionic acid are dissolved in 4 ml of 1N NaOH + 100 ml of water. The solution is freeze-dried, and the sodium salt of the carboxylic acid used is obtained quantitatively.

10 10 g (34.9 mmol) of methyl 2-hydroxy-3-methoxy-3,3-diphenylpropionate were dissolved in 50 ml each of methanol and glacial acetic acid, 1 ml of RuO(OH)₂ in dioxane was added, and hydrogenation was carried out with H₂ in an autoclave at 100°C under 100 bar for 30 h. The catalyst was filtered off, the mixture was concentrated, mixed with ether and washed with NaCl solution, and the organic phase was dried and concentrated. 10.1 g of methyl 3,3-dicyclohexyl-2-hydroxy-3-methoxypropionate were obtained as an oil.

25 Example 7

Methyl 2-[(4,6-dimethoxy-pyrimidin-2-yl)thio]-3-methoxy-3,3-diphenylpropionate [sic]

7.16 g (25 mmol) of methyl 2-hydroxy-3-methoxy-3,3-diphenylpropionate were dissolved in 50 ml of dichloromethane, 3 g (30 mmol) of triethylamine were added, and 3.2 g (28 mmol) of methanesulfonyl chloride were added dropwise while stirring. The mixture was stirred at room temperature for 2 h, washed with water, dried over magnesium sulfate and concentrated under reduced pressure. The residue was taken up in DMF and added dropwise at 0°C to a suspension of 12.9 g (75 mmol) of 4,6-dimethoxypyrimidine-2-thiol and 8.4 g (100 mmol) of sodium bicarbonate in 100 ml of DMF. After stirring at room temperature for 2 h and at 60°C for a further 2 h, the mixture was poured into 1 liter of ice-water, and the resulting precipitate was filtered off with suction. After drying, 3.19 g (29 %) of a white powder remained.

Example 8

Methyl 2-hydroxy-3,3-diphenylbutyrate

1.5 g (5.9 mmol) of methyl 3,3-diphenyl-2,3-epoxypropionate dissolved in 10 ml of absolute ether were added dropwise to a cuprate solution which had been prepared from 635 mg (7 mmol) of copper(I) cyanide dissolved in 10 ml of absolute ether and 8.14 ml (13 mmol) of a 1.6 normal methyllithium solution and had been cooled to -78°C . The solution was stirred at -78°C for 1 h and then allowed to warm to room temperature. It was subsequently diluted with 100 ml of ether and 100 ml of water, and the ether phase was washed with dilute citric acid and with sodium bicarbonate solution and dried over magnesium sulfate. The crude product was purified by chromatography on silica gel with cyclohexane/ethyl acetate mixtures to result in 250 mg (16 %) of a pale yellow oil.

Example 9

2-Hydroxy-3-methoxy-3,3-diphenylpropionic acid

91.11 g (0.5 mol) of benzophenone and 45.92 g (0.85 mol) of sodium methoxide were suspended in 150 ml of methyl tert-butyl ether (MTB) at room temperature. After cooling to -10°C , 92.24 g (0.85 mol) of methyl chloroacetate were added in such a way that the internal temperature rose to 40°C while continuing to cool in a bath at -10°C . The mixture was then stirred without cooling at the autogenous temperature for one hour. After addition of 250 ml of water and brief stirring, the aqueous phase was separated off. The MTB phase was washed with 250 ml of dilute sodium chloride solution. After the solvent had been changed to methanol (250 ml), a solution of 1 g of p-toluenesulfonic acid in 10 ml of methanol was added at room temperature. The mixture was stirred at autogenous temperature for one hour and then heated to reflux. While distilling out the methanol, 400 g of a 10 % strength sodium hydroxide solution was added dropwise, and finally 60 ml of water were added. The methanol was distilled out until the bottom temperature reached 97°C . After cooling to 55°C , 190 ml of MTB were added and the mixture was acidified to pH 2 with about 77 ml of concentrated HCl. After cooling to room temperature, the aqueous phase was separated off and the organic phase was concentrated by distilling out 60 ml of MtB [sic]. The product was crystallized by adding 500 ml of heptane and slowly cooling to room temperature. The coarsely crystalline solid was filtered off with suction, washed with heptane and dried to constant weight in a vacuum oven at 40°C .

Yield: 108.9 g (80 %), HPLC > 99.5 % area.

Example 10

S-2-Hydroxy-3-methoxy-3,3-diphenylpropionic acid (racemate resolution with L-proline methyl ester)

5 148.8 g of a 30 % strength methanolic sodium methanolate solution (0.826 mol) were added dropwise to 240 g of a 57 % strength methanolic L-proline methyl ester hydrochloride solution (0.826 mol) at room temperature, and 2.4 l of MTB and 225 g (0.826 mol) of 2-hydroxy-3-methoxy-3,3-diphenylpropionic acid
10 were added. After 2680 ml of MTB/methanol mixture had been distilled out with simultaneous dropwise addition of 2.4 l of MTB, the mixture was slowly cooled to room temperature, the crystals (R-2-hydroxy-3-methoxy-3,3-diphenylpropionic acid x L-proline methyl ester) were filtered off with suction, and the solid was
15 washed with 150 ml of MTB. The filtrate was concentrated by distilling out 1.5 l of MTB, and 1.0 l of water was added. The pH was adjusted to 1.2 with concentrated hydrochloric acid at room temperature and, after stirring and phase separation, the aqueous phase was separated off and extracted with 0.4 l of MTB. The com-
20 bined organic phases were extracted with 0.4 l of water. The residue after the MTB had been stripped off was dissolved in 650 ml of toluene under reflux, and the product was crystallized by seeding and slow cooling. Filtration with suction, washing with toluene and drying in a vacuum oven resulted in 78.7 g of
25 S-2-hydroxy-3-methoxy-3,3-diphenylpropionic acid (yield 35 % based on the racemate).
Chiral HPLC: 100 % pure
HPLC: 99.8 %

30 Example 11

S-2-Hydroxy-3-methoxy-3,3-diphenylpropionic acid (racemate resolution with (S)-1-(4-nitrophenyl)ethylamine)

30.5 g (0.184 mol) of (S)-1-(4-nitrophenyl)ethylamine were added
35 to 100 g (0.368 mol) of 2-hydroxy-3-methoxy-3,3-diphenylpropionic acid in 750 ml of acetone and 750 ml of MTB under reflux, the mixture was seeded, boiled under reflux for one hour and slowly cooled to room temperature for crystallization. The crystals (S-2-hydroxy-3-methoxy-3,3-diphenylpropionic acid x (S)-1-(4-ni-
40 trophenyl)ethylamine) were filtered off with suction and washed with MTB. The residue was suspended in 500 ml of water and 350 ml of MTB and then the pH was adjusted to 1.2 with concentrated hydrochloric acid at room temperature, and, after stirring and phase separation, the aqueous phase was separated off and ex-
45 tracted with 150 ml of MTB. The combined organic phases were extracted with 100 ml of water. 370 ml of MTB were distilled out and then 390 ml of n-heptane were added under reflux, and the

mixture was slowly cooled to room temperature while the product crystallized. Filtration with suction, washing with n-heptane and drying in a vacuum oven resulted in 35.0 g of S-2-hydroxy-3-methoxy-3,3-diphenylpropionic acid (yield 35 % based on the racemate).

Chiral HPLC: 100 % pure

HPLC: 99.8 %

Example 12

10 Benzyl 3-methoxy-2-(4-methoxy-6,7-dihydro-5H-cyclopentapyrimidin-2-yloxy)-3,3-diphenylpropionate

24.48 g (90 mmol) of 3-methoxy-3,3-diphenyl-2-hydroxypropionic acid were dissolved in 150 ml of DMF, and 13.7 g (99 mmol) of potassium carbonate were added. The suspension was stirred at room temperature for 30 min. Then 10.7 ml (90 mmol) of benzyl bromide were added dropwise over the course of 5 min, and the mixture was stirred for 1 h, during which the temperature rose to 32°C.

20

To this mixture were successively added 24.84 g (180 mmol) of K_2CO_3 and 20.52 g (90 mmol) of 2-methanesulfonyl-4-methoxy-6,7-dihydro-5H-cyclopentapyridine [sic], and the mixture was stirred at 80°C for 3 h.

25

For workup, the contents of the flask were diluted with about 600 ml of H_2O and cautiously acidified with concentrated HCl, and 250 ml of ethyl acetate were added. 31.4 g of pure product precipitated and were filtered off.

30

The ethyl acetate phase was separated from the mother liquor, the aqueous phase was extracted again with ethyl acetate, and the combined organic phases were concentrated. The oily residue (19 g) was purified by chromatography (cyclohexane/ethyl acetate = 9/1) to result in a further 10.5 g of pure product.

35

Total yield: 41.9 g (82.2 mmol) \cong 91 %

Melting point 143-147°C

MS: MH^+ = 511

40

Example 13

3-Methoxy-2-(4-methoxy-(6,7-dihydro-5H-cyclopentapyrimidin-2-yloxy)-3,3-diphenylpropionic [sic] acid

45 40 g (78.4 mmol) of benzyl 3-methoxy-2-(4-methoxy-6,7-dihydro-5H-cyclopentapyrimidin-2-yloxy)-3,3-diphenylpropionate were dissolved in 400 ml of ethyl acetate/methanol (4:1), about 500 mg

of palladium on active carbon (10 %) were added, and the mixture was exposed to a hydrogen atmosphere until no further gas was taken up. The catalyst was filtered off, the solution was evaporated, and the residue was crystallized from ether.

5

Example 14

Ethyl 2S-3,3-diphenyloxirane-2-carboxylate

2.57 g (10.2 mmol) of ethyl 3,3-diphenylacrylate and 464 mg of
10 4-phenylpyridine N-oxide were dissolved in 24 ml of methylene
chloride, and 432 mg (6.5 mol%) of (5,5)-(+) -N,N'-bis(3,5-di-
tert-butylsalicylidene)-1,2-cyclohexanediaminomanganese(III)
chloride were added. While cooling in ice, 6.4 ml of a 12 %
strength sodium hypochloride [sic] solution were added, and the
15 mixture was stirred while cooling in ice for 30 min and at room
temperature overnight. The solution was diluted to 200 ml with
water, extracted with ether, dried and evaporated. 2.85 g of a
colorless oil were obtained. Purification by NPLC [sic] (cyclo-
hexane:ethyl acetate = 9:1) resulted in 1.12 g of oil with an
20 enantiomer ratio of about 8:1 in favor of the S configuration.

¹H-NMR [CDCl₃],

δ = 1.0 (t, 3H); 3.9 (m, 3H); 7.3 (m, 10H)

25 Example 15

2-Methylsulfonyl-6,7-dihydro-5H-cyclopentapyrimidin-4-ol [sic]

46.9 g (330 mmol) of methyl cyclopentanone-2-carboxylate and
53.5 g (192 mmol) of 5-methylisothiourea [sic] sulfate were suc-
30 cessively added to 29.6 g (528 mmol) of KOH in 396 ml of meth-
anol, and the mixture was stirred at room temperature overnight,
acidified with 1N hydrochloric acid and diluted with water. The
crystals which separated out were filtered off with suction and
dried. 20 g of crystals were obtained.

35

Example 16

sulfanyl 4-Chloro-2-methyl-6,7-dihydro-5H-cyclopentapyrimidine
[sic]

40 255 ml of phosphorus oxychloride were added to 20 g (110 mmol)
[lacuna], and the mixture was stirred at 80°C for 3 hours. Phos-
phorus oxychloride was evaporated off, ice was added to the resi-
due, and the crystals which separated out were filtered off with
suction. 18.5 g of a brownish solid were obtained.

45

Example 17

4-Methoxy-2-methylsulfonyl-6,7-dihydro-5H-cyclopentapyrimidine
[sic]

5 18.05 g (90 mmol) of 4-chloro-2-methylsulfonyl-6,7-dihydro-5H-cyclopentapyrimidine [sic] were dissolved in 200 ml of methanol. At 45°C, 16.7 g of sodium methoxide (as 30 % strength solutions [sic] in methanol) were added dropwise, and the mixture was stirred for 2 hours. The solution was evaporated, taken up in
10 ethyl acetate and acidified with dilute hydrochloric acid, and the ethyl acetate extract was evaporated. 15.5 g of an oil remained.

¹H-NMR [DMSO],

15 δ = 2.1 (quintet, 2H); 2.5 (s, 3H);
2.8 (dt, 4H); 3.9 (s, 3H) ppm

Example 18

2-Methylsulfonyl-4-methoxy-6,7-dihydro-5H-cyclopentopyrimidine
20 [sic]

15 g (76.2 mmol) of 4-methoxy-2-methylsulfonyl-6,7-dihydro-5H-cyclopentapyrimidine [sic] were dissolved in 160 ml of glacial acetic acid/methylene chloride (1:1), and 1.3 g of sodium tungstate were added. At 35°C, 17.5 ml (170 ml [sic]) of a 30 %
25 strength H₂O₂ solution were added dropwise. The mixture was then diluted with 500 ml of water and 100 ml of methylene chloride, and the organic phase was separated off, dried and evaporated. 14 g of oil remained and were crystallized from ether.

30

¹H-NMR [CDCl₃],

δ = 2.2 (quintet, 2H); 3.0 (dt., 4H);
3.3 (s, 3H); 4.1 (s, 3H) ppm

35 Example 19

1-Benzenesulfonyl-3-(4,6-dimethoxy-2-pyrimidinyl-4-methoxy-4,4-diphenyl-2-butanone

0.37 g (2.4 mmol) of phenyl methane [sic] sulfone were dissolved
40 in 10 ml of dry THF and then, at -70°C, 2 eq. of butyllithium (2.94 ml; 1.6 molar solution in hexane) were added dropwise. After 1 h at -70°C, 1 g (2.4 mmol) of methyl 2-(4,6-dimethoxy-2-pyrimidinyl-3-methoxy-3,3-diphenylpropynoate. [sic]
dissolved in 5 ml of THF was added dropwise. The reaction mixture
45 was then stirred at -70°C for 1 h and at -10°C for 1 h and then warmed to room temperature.

For workup, about 10 ml of saturated NH_4Cl solution were added dropwise, thorough extraction with ethyl acetate was carried out, and the combined organic phases [lacuna] with saturated N-Cl [sic] solution and dried over Na_2SO_4 . The residue obtained after 5 drying and concentration was purified by chromatography on silica gel (n -heptane/ethyl acetate 15 % \rightarrow 30 %) and subsequently MPLC on RP silica gel (acetonitrile/ H_2O + TFA); 0.3 g of a white amorphous powder was obtained as product.

10 Example 20

3,3-Diphenyloxiram-2-carbonitrile [sic]

3.1 g (54.9 mmol) of sodium methoxide were suspended in 20 ml of dry THF and then, at -10°C , a mixture of 5 g (27.4 mmol) of benzo-15 phenone and 4.2 g (54.9 mmol) of chloroacetonitrile was added dropwise.

The reaction mixture was stirred at -10°C for about 2 h, then poured into water and extracted several times with ethyl acetate.

20 The combined organic phases were dried over Na_2SO_4 and concentrated, and the residue was purified by chromatography on silica gel (n -heptane/ethyl acetate).

Yield: 1.2 g (20 %)

25 $^1\text{H-NMR}$ [CDCl_3],

δ = 3.9 (s, 1H); 7.4-7.5 (m, 10 H) ppm

Example 21

2-Hydroxy-3-methoxy-3,3-diphenylpropionitrile

30

6.5 [lacuna] (29.4 mmol) of 3,3-diphenyloxirane-2-carbonitrile were dissolved in 60 ml of methanol and, at 0°C , about 2 ml of boron trifluoride etherate solution were added. The mixture was stirred further at 0°C for 1 h and then at room temperature over-35 night. For workup it was diluted with diethyl ether and washed with saturated NaCl solution, and the organic phase was dried over Na_2SO_4 and concentrated. The residue comprised 7.3 g of a white amorphous powder which was used directly in the subsequent reactions.

40

$^1\text{H-NMR}$ [CDCl_3],

δ = 2.95 (broad s, OH), 3.15 (s, 3H),

5.3 (s, 1H), 7.3-7.5 (m, 10) ppm

45

Example 22

2-(4,6-Dimethoxy-2-pyrimidinyloxy)-3-methoxy-3,3-diphenylpropionitrile

5 7.3 g (28.8 mmol) of 2-hydroxy-3-methoxy-3,3-diphenylpropionitrile were dissolved in 90 ml of DMF, and 4 g (28.8 mmol) of K_2CO_3 and 6.3 g (28 mmol) of 2-methanesulfonyl-4,6-dimethoxypyrimidine were added. The mixture was stirred at room temperature for about 12 h, then poured into water and extracted with ethyl
10 acetate. The combined organic phases were washed again with H_2O , dried and concentrated. The residue obtained in this way was then purified by chromatography on silica gel (n-heptane/ethyl acetate).

Yield: 6.9 g of white amorphous powder

15

FAB-MS: 392 ($M+H^+$)

1H -NMR [$CDCl_3$],

δ = 3.3 (s, 3H); 4.95 (s, 6H), 5.85 (s, 1H);

6.3 (s, 1H); 7.3-7.5 (m, 10H) ppm

20

Example 23

5-[2-(4,6-Dimethoxy-2-pyrimidinyloxy)-3-methoxy-3,3-diphenylpropyl]-1H-tetrazole [sic]

25 0.5 g (1.3 mmol) of nitrile was dissolved in 10 ml of toluene, and 85 mg (1.3 mmol) of NaN_3 and 460 mg (1.4 mmol) of Bu_3SnCl were successively added, and then the mixture was refluxed for about 40 h. Cooling was followed by dilution with ethyl acetate and washing with 10 % aqueous KF solution and with NaCl solution.
30 After drying over $MgSO_4$ and concentration there remained 1.0 g of a yellow oil, which was purified by chromatography on silica gel (n-heptane/ethyl acetate).

Concentration of the fractions resulted in 60 mg of the 1H-tetrazole and 110 mg of the 1-methyltetrazole, each as amorphous white solids.
35

5-[2-(4,6-Dimethoxy-2-pyrimidinyloxy)-3-methoxy-3,3-diphenylpropyl]-1H-tetrazole [sic]

40

Electrospray-MS: 435 ($M+H^+$)

1H -NMR ($CDCl_3$):

δ (ppm) 3.28 (s, 3H), 3.85 (s, 6H), 5.75 (s, 1H), 7.25-7.40 (m, 10H), 7.50 (s, 1H).
45

5-[2-(4,6-Dimethoxy-2-pyrimidinyloxy)-3-methoxy-3,3-diphenyl)-propyl]-1-methyltetrazole [sic]

Electrospray-MS; 471 (M+H⁺)

5 ¹H-NMR (CDCl₃):

δ (ppm) 3.0 (s, 3H), 3.35 (s, 3H) [sic], 3.80 (s, 6H), 5.75 (s, 1H), 7.30-7.40 (m, 11H).

Example 24

10 2-(4,6-Dimethoxy-2-pyrimidinyloxy)-3-methylsulfinyl-3,3-diphenylpropionic acid

1.2 g (2.9 mmol) of

2-(4,6-dimethoxy-2-pyrimidinyloxy)-3-methylsulfonyl-3,3-diphenyl-
15 propionic [sic] acid were introduced into 15 ml of glacial acetic acid at 0°C and 294 μl of 30 % strength H₂O₂ were added dropwise. The mixture was stirred at room temperature overnight, poured into water, extracted with CH₂Cl₂ and washed with sodium thiosulfate solution and brine. After drying, 1 g of substance was iso-
20 lated as a white foam.

Example 25

2-(4,6-Dimethoxy-2-pyrimidinyloxy)-3-methylsulfonyl-3,3-diphenylpropionic acid

25


0.6 g (1.45 mmol) of 2-(4,6-dimethoxy-2-pyrimidinyloxy)-3-methylsulfonyl-3,3-diphenylpropionic [sic] acid was introduced into 15 ml of glacial acetic acid at room temperature, and 294 μl of 30 % strength H₂O₂ were added dropwise. The mixture was stirred at room
30 temperature overnight, heated at 50°C for a further 3 h, poured into water and washed with sodium thiosulfate solution and brine. After drying, 400 mg were isolated as a white solid.

The compounds listed in Table 1 [sic] can be prepared in a similar way.
35

40

45

Table I



No.	R ¹	R ⁴ , R ⁵	R ⁶	R ²	R ³	X	Y	Z	m.p.[°C]
1-195	OMe	Phenyl	Methyl	OMe	OMe	CH	O	O	81
1-196	OH	Phenyl	Methyl	OMe	OMe	CH	O	O	167
1-197	OH	Phenyl	CH ₂ -CH ₂ -S-CH ₃	OMe	OMe	CH	O	O	
1-198	OH	Phenyl	Ethyl	OMe	OMe	CH	O	O	81 (decomp.)
1-199	OH	Phenyl	iso-Propyl	OMe	OMe	CH	O	O	182
1-200	OH	Phenyl	Methyl	OMe	OMe	CH	O	S	168
1-201	OH	Phenyl	CH ₂ -CH ₂ -SO ₂ -CH(CH ₃) ₂	OMe	OMe	CH	O	O	
1-202	OH	Phenyl	CH ₂ -CH ₂ -SO ₂ -CH(CH ₃) ₂	OMe	OMe	CH	S	O	
1-203	OH	Phenyl	CH ₂ -CH ₂ -SO ₂ -CH(CH ₃) ₂	OMe	OMe	C-CH(CH ₃) ₂	O	O	
1-204	OH	Phenyl	CH ₂ -CH ₂ -SO ₂ -CH(CH ₃) ₂	OMe	OMe	C-CH(CH ₃) ₃	O	O	
1-205	OH	Phenyl	CH ₂ -CH ₂ -SO ₂ -CH(CH ₃) ₂	OMe	NH-OCH ₃	CH	O	O	
1-206	OH	Phenyl	n-Propyl	OMe	OMe	CH	O	O	174
1-207	OMe	Phenyl	n-Propyl	OMe	OMe	CH	O	O	
1-208	OH	Phenyl	n-Propyl	OEt	OEt	CH	O	O	

No.	R ¹	R ⁴ , R ⁵	R ⁶	R ²	R ³	X	Y	Z	m.p.[°C]
I-209	OH	Phenyl	n-Butyl	OMe	OMe	CH	O	O	
I-210	OH	Phenyl	iso-Butyl	OMe	OMe	CH	O	O	
I-211	OH	Phenyl	iso-Butyl	OMe	O-CH ₂ -CH ₂ -C		O	O	
I-212	OH	Phenyl	tert-Butyl	OMe	OMe	CH	O	O	
I-213	OH	Phenyl	Cyclopropyl	OMe	OMe	CH	O	O	
I-214	OH	Phenyl	Cyclopentyl	OMe	OMe	CH	O	O	
I-215	OH	Phenyl	Cyclohexyl	OMe	OMe	CH	O	O	
I-216	OH	Phenyl	(CH ₃) ₃ C-CH ₂ -CH ₂	OMe	OMe	CH	O	O	
I-217	OH	Phenyl	(CH ₃) ₂ CH-CH ₂ -CH ₂ -CH ₂	OMe	OMe	CH	O	O	173
I-218	OH	Phenyl	HO-CH ₂ -CH ₂	OMe	OMe	CH	O	O	
I-219	OH	Phenyl	HO ₂ C-(CH ₂) ₂ -	OMe	OMe	CH	O	O	
I-220	OH	Phenyl	Cyclopropylmethylene [sic]	OMe	OMe	CH	O	O	115
I-221	OH	Phenyl	H	OMe	OMe	CH	O	O	
I-222	OH	Phenyl	Methyl	OMe	OMe	CH	O	O	
I-223	OH	Phenyl	Phenyl	OMe	OMe	CH	O	O	136
I-224	OH	Phenyl	Phenyl	OMe	O-CH(CH ₃)-CH ₂ -C		O	O	
I-225	OMe	Phenyl	Phenyl	OMe	OMe	CH	O	O	
I-226	OH	Phenyl	4-Isopropyl-Phenyl	OMe	OMe	CH	O	O	
I-227	OH	Phenyl	4-Me-S-Phenyl	OMe	OMe	CH	O	O	
I-228	OH	Phenyl	4-Me-O-Phenyl	OMe	OMe	CH	O	O	
I-229	OH	Phenyl	3-Et-Phenyl	OMe	OMe	CH	O	O	
I-230	OH	Phenyl	2-Me-Phenyl	OMe	OMe	CH	O	O	

No.	R ¹	R ⁴ , R ⁵	R ⁶	R ²	R ³	X	Y	Z	m.p. [°C]
I-231	OH	Phenyl	2-Cl-Phenyl	OMe	OMe	CH	O	O	
I-232	OH	Phenyl	3-Br-Phenyl	OMe	OMe	CH	O	O	
I-233	OH	Phenyl	4-F-Phenyl	OMe	OMe	CH	O	O	
I-234	OH	Phenyl	4-F-Phenyl	OMe	OMe	CH	S	O	
I-235	OH	Phenyl	4-CH ₃ -Phenyl	OMe	OMe	CH	O	O	
I-236	OH	Phenyl	3-NO ₂ -Phenyl	OMe	OMe	CH	O	O	
I-237	OH	Phenyl	2-HO-Phenyl	OMe	OMe	CH	O	O	
I-238	OH	Phenyl	3,4-Dimethoxyphenyl	OMe	OMe	CH	O	O	
I-239	OH	Phenyl	3,4-Dioxomethylenepheryl [sic]	OMe	OMe	CH	O	O	
I-240	OH	Phenyl	3,4,5-Trimethoxyphenyl	OMe	OMe	CH	O	O	
I-241	OH	Phenyl	Benzyl	OMe	OMe	CH	O	O	
I-242	OH	Phenyl	2-Cl-Benzyl	OMe	OMe	CH	O	O	
I-243	OH	Phenyl	3-Br-Benzyl	OMe	OMe	CH	O	O	
I-244	OH	Phenyl	4-F-Benzyl	OMe	OMe	CH	O	O	
I-245	OH	Phenyl	2-Me-Benzyl	OMe	OMe	CH	O	O	
I-246	OH	Phenyl	2-Me-Benzyl	OMe	OMe	O-CH=CH-C	O	O	
I-247	OH	Phenyl	3-Et-Benzyl	OMe	OMe	CH	O	O	
I-248	OH	Phenyl	4-iso-Propyl-Benzyl	OMe	OMe	CH	O	O	
I-249	OH	Phenyl	4-NO ₂ -Propyl-Benzyl	OMe	OMe	CH	O	O	
I-250	OH	Phenyl	2-Me-5-Propyl-Benzyl	OMe	OMe	CH	O	O	
I-251	OH	Phenyl	2-Me-5-Propyl-Benzyl	OEt	OEt	CH	O	O	
I-252	OH	Phenyl	4-Me-2-Propyl-Benzyl	OMe	OMe	CH	O	O	

No.	R ¹	R ⁴ , R ⁵	R ⁶	R ²	R ³	X	Y	Z	m.p. [°C]
I-253	OH	Phenyl	3,4-Dioxomethylenebenzyl [sic]	OMe	OMe	CH	O	O	
I-254	OH	4-F-Phenyl	Methyl	OMe	OMe	CH	O	O	163-165 (decomp.)
I-255	OMe	4-F-Phenyl	Methyl	OEt	OEt	CH	O	O	
I-256	OH	4-Cl-Phenyl	Methyl	OMe	OMe	CH	O	O	
I-257	OH	4-Me-O-Phenyl	Methyl	OMe	OMe	CH	O	O	
I-258	OH	4-Me-O-Phenyl	Ethyl	OMe	OMe	CH	O	O	
I-259	OH	4-Me-Phenyl	Methyl	OMe	OMe	CH	O	O	
I-260	OH	4-Me-Phenyl	Methyl	OMe	OMe	O-CH ₂ -CH ₂ -C	O	O	
I-261	OH	3-CF ₃ -Phenyl	n-Propyl	OMe	OMe	CH	O	O	
I-262	OH	3-CF ₃ -Phenyl	n-Propyl	OMe	OMe	O-CH(CH ₃)-CH ₂ -C	O	O	
I-263	OH	4-NO ₂ -Phenyl	Methyl	OMe	OMe	CH	O	O	
I-264	OH	4-NO ₂ -Phenyl	Methyl	OMe	OMe	O-CH=CH-C	O	O	
I-265	OH	3-Cl-Phenyl	Ethyl	OMe	OMe	CH	O	O	
I-266	OH	2-F-Phenyl	Methyl	OMe	OMe	CH	O	O	193-194 (decomp.)
I-267	OH	2-F-Phenyl	Methyl	OMe	OMe	CH	S	O	
I-268	OH	2-Me-O-Phenyl	Methyl	OMe	OMe	CH	O	O	
I-269	OH	2-Me-O-Phenyl	Methyl	OMe	OMe	CH	O	S	
I-270	OH	3,4-Dimethoxyphenyl	Methyl	OMe	OMe	CH	O	O	
I-271	OH	3,4-Dioxomethylenepheryl [sic]	Methyl	OMe	OMe	CH	O	O	
I-272	OH	p-CF ₃ -Phenyl	Methyl	OMe	OMe	CH	O	O	

59

60

65

70

75

78

No.	R ¹	R ⁴ , R ⁵	R ⁶	R ²	R ³	X	Y	Z	m.p.[°C]
I-273	OH	Phenyl	Methyl	OMe	OEt	CH	O	O	
I-274	OMe	Phenyl	Methyl	OMe	OEt	CH	S	O	
I-275	OH	Phenyl	Ethyl	OMe	NH-OMe	CH	O	O	
I-276	OH	p-Me-O-Phenyl	n-Propyl	OMe	OCF ₃	CH	O	O	
I-277	OH	Phenyl	Methyl	OMe	CF ₃	CH	O	O	
I-278	OH	Phenyl	Methyl	OMe	CF ₃	N	O	O	
I-279	OH	3,4-Dimethoxyphenyl	Benzyl	Me	Me		O	O	
I-280	OH	3,4-Dimethoxyphenyl	Methyl	OMe	OMe	O-CH ₂ -CH ₂ -C	O	O	
I-281	OH	Phenyl	Methyl	OMe	OMe	O-CH ₂ -CH ₂ -C	O	O	126 (decomp.)
I-282	OH	Phenyl	Methyl	OMe	OMe	O-CH(CH ₃)-CH ₂ -C	O	O	
I-283	OH	Phenyl	Methyl	OMe	OMe	N(CH ₃)-CH=CH-C	O	O	118
I-284	OH	Phenyl	Methyl	OMe	OMe	S-C(CH ₃)=C(CH ₃)-C	O	O	
I-285	OH	Phenyl	Methyl	OMe	OMe	O-C(CH ₃)=CH-C	O	O	
I-286	OH	Phenyl	Methyl	Me	Me	O-C(CH ₃)=CH-C	O	O	
I-287	OH	Phenyl	Methyl	Me	Me	O-CH=CH-C	O	O	
I-288	OH	4-F-phenyl	Methyl	Me	Me	S-CH=CH-C	O	O	
I-289	OH	4-F-phenyl	H	OMe	OMe	CH	O	O	
I-290	OH	Phenyl	Methyl	OMe	OMe	CH ₂ -CH ₂ -CH ₂ -C	O	O	149-151 (decomp.)
I-291	OH	Phenyl	Methyl	Methyl	Methyl	CH ₂ -CH ₂ -CH ₂ -C	O	O	157 (decomp.)
I-292	OH	Phenyl	Methyl	Ethyl	Ethyl	CH ₂ -CH ₂ -CH ₂ -CH ₂ -C	O	O	
I-293	OH	Phenyl	Methyl	OMe	OMe	CH ₂ -CH ₂ -CH ₂ -CH ₂ -C	O	O	
I-294	OH	Phenyl	Methyl	Me	Me	CH	O	O	

No.	R ¹	R ⁴ , R ⁵	R ⁶	R ²	R ³	X	Y	Z	m.p. [°C]
I-295	OH	Phenyl	Methyl	Et	Et	CH	O	O	
I-296	OH	Phenyl	Methyl	Me	Me	C-CH ₃	O	O	
I-297	OH	Phenyl	Methyl	OMe	Me	CH	O	O	
I-298	OH	Cyclohexyl	Methyl	OMe	OMe	CH	O	O	
I-299	OH	Cyclohexyl	Methyl	OMe	CH ₂ -CH ₂ -CH ₂ -C		O	O	
I-300	OH	Phenyl	Methyl	OCH ₃	OCH ₃	CH	S	S	
I-301	OH	Phenyl	Methyl	OCH ₃	OCH ₃	CH	O	S	134
I-302	OCH ₃	Phenyl	Methyl	OCH ₃	OCH ₃	CH	S	S	
I-303	OH	Phenyl	Methyl	OCH ₃	OCH ₃	CH	O	O	
I-304	OCH ₃	2-Fluorophenyl	Methyl	OCH ₃	OCH ₃	CH	O	O	
I-305	OC ₂ H ₅	3-Chlorophenyl	Methyl	OCH ₃	OCH ₃	N	O	O	
I-306	ON(CH ₃) ₂	4-Bromophenyl	Methyl	CF ₃	CF ₃	CH	S	O	
I-307	O-CH ₂ -C=CH	Phenyl	Ethyl	OCH ₃	CF ₃	CH	O	O	
I-308	OH	Phenyl	Propyl	OCH ₃	OCF ₃	CH	O	S	
I-309	OCH ₃	Phenyl	i-Propyl	OCH ₃	CH ₃	CH	O	O	
I-310	OC ₂ H ₅	Phenyl	s-Butyl	OCH ₃	Cl	CH	S	O	
I-311	ON(CH ₃) ₂	2-Methylphenyl	Methyl	OCH ₃	OCH ₃	CH	O	O	
I-312	ON(CH ₃) ₂	3-Methoxyphenyl	Methyl	OCH ₃	OCH ₃	CH	O	O	
I-313	ON=C(CH ₃) ₂	4-Nitrophenyl	Methyl	OCH ₃	OCH ₃	CH	O	O	
I-314	ON(CH ₃) ₂	Phenyl	1-Phenylpropyn-3-yl	OCH ₃	OCF ₃	N	O	S	
I-315	ON=C(CH ₃) ₂	2-Hydroxyphenyl	Methyl	OCH ₃	CH ₃	N	O	O	
I-316	ONSO ₂ C ₆ H ₅	3-Trifluoromethylphenyl	Methyl	OCH ₃	Cl	N	O	O	

No.	R ¹	R ⁴ , R ⁵	R ⁶	R ²	R ³	X	Y	Z	m.p.[°C]
I-317	NHPhenyl	4-Dimethylaminophenyl	Methyl	OCH ₃	OCH ₃	CH	S	O	
I-318	OC ₂ H ₅	Phenyl	Trifluoroethyl	CH ₃	CH ₃	CH	O	O	
I-319	ON(CH ₃) ₂	Phenyl	Benzyl	Cl	Cl	CH	O	O	
I-320	ON(CH ₃) ₂	Phenyl	2-Methoxyethyl	OCH ₃	-O-CH ₂ -CH ₂ -		S	O	
I-321	OH	Phenyl	Phenyl	OCH ₃	OCH ₃	CH	O	O	
I-322	OH	Phenyl	Phenyl	OCH ₃	-O-CH ₂ -CH ₂ -		O	O	
I-323	OH	Phenyl	Phenyl	OCH ₃	OCH ₃	N	O	O	
I-324	OH	Phenyl	Phenyl	OCH ₃	OCH ₃	CH	S	O	
I-325	OH	Phenyl	Phenyl	OCH ₃	OCH ₃	CH	S	S	
I-326	OH	Phenyl	Phenyl	OCH ₃	OCH ₃	CH	O	S	
I-327	OH	Phenyl	Phenyl	OCH ₃	OCH ₃	CH	O	O	
I-328	OH	Phenyl	Phenyl	OCH ₃	OCH ₃	CH	O	O	
I-329	OH	-(CH ₂) ₅	Phenyl	OCH ₃	OCH ₃	CH	O	O	
I-330	OH	Phenyl	2-Thiazolyl	OCH ₃	OCH ₃	CH	O	O	
I-331	OCH ₃	2-Fluorophenyl	Phenyl	OCH ₃	OCH ₃	CH	O	O	
I-332	OC ₂ H ₅	3-Chlorophenyl	Phenyl	OCH ₃	OCH ₃	CH	O	O	
I-333	ON(CH ₃) ₂	4-Bromophenyl	Phenyl	CF ₃	CF ₃	N	O	O	
I-334	O-CH ₂ ≡CH	Phenyl	2-Fluorophenyl	OCH ₃	CF ₃	CH	O	O	
I-335	OH	Phenyl	3-Chlorophenyl	OCH ₃	OCF ₃	CH	O	S	
I-336	OCH ₃	Phenyl	4-Bromophenyl	OCH ₃	CH ₃	CH	O	O	
I-337	OC ₂ H ₅	Phenyl	4-Thiazolyl	OCH ₃	Cl	CH	S	O	
I-338	ON(CH ₃) ₂	2-Methylphenyl	Phenyl	OCH ₃	OCH ₃	CH	O	O	

No.	R ¹	R ⁴ , R ⁵	R ⁶	R ²	R ³	X	Y	Z	m.p. [°C]
I-339	ON=C(CH ₃) ₂	3-Methoxyphenyl	Phenyl	OCH ₃	OCH ₃	CH	O	O	
I-340	OH	Phenyl	Methyl	OCH ₃	-CH ₂ -CH ₂ -CH ₂ -C		O	O	
I-341	OH	4-Fluorophenyl	Methyl	OCH ₃	CH		O	O	168 (decomp.)
I-342	OH	4-Fluorophenyl	Methyl	OCH ₃	-CH ₂ -CH ₂ -CH ₂ -C		O	O	
I-343	NH-SO ₂ -C ₆ H ₅	4-Nitrophenyl	Phenyl	OCH ₃	OCH ₃	CH	O	O	
I-344	OCH ₃	Phenyl	3-Imidazolyl	OCH ₃	-O-CH ₂ -CH ₂		O	O	
I-345	OC ₂ H ₅	Phenyl	4-Imidazolyl	OCH ₃	CF ₃	N	S	O	
I-346	ON(CH ₃) ₂	Phenyl	2-Pyrazolyl	OCH ₃	OCF ₃	N	O	S	
I-347	ON=C(CH ₃) ₂	2-Hydroxyphenyl	Phenyl	OCH ₃	CH ₃	N	O	O	
I-348	NH-SO ₂ -C ₆ H ₅	3-Trifluoromethylphenyl	Phenyl	OCH ₃	Cl	N	O	O	
I-349	NHPhenyl	4-Dimethylaminophenyl	Phenyl	OCH ₃	OCH ₃	CH	S	O	
I-350	ONa	Phenyl	Phenyl	OCH ₃	OCH ₃	CH	S	S	
I-351	O-CH ₂ -C≡C	Phenyl	Phenyl	OCH ₃	OCH ₃	N	S	S	
I-352	OH	Phenyl	Phenyl	CF ₃	CF ₃	CH	O	S	
I-353	OCH ₃	Phenyl	Phenyl	OCF ₃	OCF ₃	CH	O	O	
I-354	OC ₂ H ₅	Phenyl	2-Dimethylaminophenyl	CH ₃	CH ₃	CH	O	O	
I-355	ON(CH ₃) ₂	Phenyl	3-Hydroxyphenyl	Cl	Cl	CH	O	O	
I-356	ON=C(CH ₃) ₂	Phenyl	4-Trifluoromethylphenyl	OCH ₃	-O-CH ₂ -CH ₂ -		S	O	
I-357	NH-SO ₂ -C ₆ H ₅	Phenyl	2-Oxazolyl	OCH ₃	CF ₃	N	S	S	
I-358	OH	Phenyl	Methyl	CH ₃	CH ₃	CH	O	O	
I-359	OH	Cyclohexyl	Methyl	OCH ₃	OCH ₃	CH	O	O	
I-360	OH	Cyclohexyl	Methyl	OCH ₃	CH ₂ -CH ₂ -CH ₂ -C		O	O	

No.	R ¹	R ⁴ , R ⁵	R ⁶	R ²	R ³	X	Y	Z	m.p. [°C]
I-361	OH	Phenyl	Methyl	N(CH ₃) ₂	N(CH ₃) ₂	CH	O	O	
I-362	OH	Phenyl	Methyl	OCH ₃	OCH ₃	CH	O	SO ₂	
I-363	OH	Phenyl	Methyl	OCH ₃	OCH ₃	CH	O	SO ₂	
I-364	OH	3-F-Phenyl	Me	OMe	OMe	CH	O	O	
I-365	OH	3-F-Phenyl	Me	OMe	CH ₂ -CH ₂ -CH ₂ -C		O	O	
I-366	OH	4-F-Phenyl	Me	OMe	CH ₂ -CH ₂ -CH ₂ -C		O	O	142-143 191°C
I-367	OH	3-MeO-Phenyl	Me	OMe	CH ₂ -CH ₂ -CH ₂ -C		O	O	158-161 (decomp.)
I-368	OH	3-MeO-Phenyl	Me	OMe	OMe	CH	O	O	
I-369	OH	3-MeO-Phenyl	Et	OMe	CH ₂ -CH ₂ -CH ₂ -C		O	O	
I-370	OH	Phenyl	HO-CH ₂ -CH ₂	OMe	CH ₂ -CH ₂ -CH ₂ -C		O	O	
I-371	OH	Phenyl	Me	NMe ₂	NMe ₂	N	O	O	181
I-372	OH	Phenyl	Me	OMe	OMe	N	O	O	
I-373	OH								
I-374	NH-SO ₂ -Phenyl	Phenyl	Me	OMe	OMe	CH	O	O	
I-375	NH-SO ₂ -Me	Phenyl	Me	OMe	OMe	CH	O	O	
I-376	CH ₂ -SO ₂ -Phenyl	Phenyl	Me	OMe	OMe	CH	O	O	
I-377	CH ₂ -SO ₂ -Me	Phenyl	Me	OMe	OMe	CH	O	O	
I-378	-CN	Phenyl	Me	OMe	OMe	CH	O	O	
I-379	Tetrazole [sic]	Phenyl	Me	OMe	OMe	CH	O	O	
I-380	NH-SO ₂ -Phenyl	Phenyl	Me	OMe	OMe	CH	O	O	167

No.	R ¹	R ⁴ , R ⁵	R ⁶	R ²	R ³	X	Y	Z	m.p. [°C]
I-381	N-Methyltetrazole [sic]	Phenyl	Me	OMe	OMe	CH	O	O	
I-382	ONa	Phenyl	Me	OMe		-O-CH ₂ -CH ₂ -C-	O	O	122-139 (zers.)
I-383	OH	o-F-Phenyl	Me	OMe		-O-CH ₂ -CH ₂ -C-	O	O	140-144 (decomp.)
I-384	OH	m-Me-Phenyl	Me	OMe	OMe	CH	O	O	169-177
I-385	OH	m-Me-Phenyl	Me	OMe		-O-CH ₂ -CH ₂ -C-	O	O	119-135 (decomp.)
I-386	OH	p-F-Phenyl	Me	OMe	Me	CH	O	O	137-140 (decomp.)
I-387	OH	m-F-Phenyl	Me	Me		-O-CH ₂ -CH ₂ -C-	O	O	150-152
I-388	OH	p-F-Phenyl	Me	Me		-O-CH ₂ -CH ₂ -C-	O	O	169-170




Table II

No.	R ¹	A	R ⁶	R ²	R ³	X	Y	Z	m.p. [°C]
II-1	OH	Bond	Methyl	OMe	OMe	CH	O	O	96-98
II-2	OH	CH ₂	Methyl	OMe	OMe	CH	O	O	

No.	R ¹	A	R ⁶	R ²	R ³	X	Y	Z	m.p.[°C]
II-3	OH	CH ₂ -CH ₂	Methyl	OMe	OMe	CH	O	O	
II-4	OH	CH=CH	Methyl	OMe	OMe	CH	O	O	
II-5	OH	O	Methyl	OMe	OMe	CH	O	O	
II-6	OH	S	Methyl	OMe	OMe	CH	O	O	
II-7	OH	NH(CH ₃)	Methyl	OMe	OMe	CH	O	O	
II-8	OH	Bond	Isopropyl	OMe	OMe	CH	O	O	137-139
II-9	OH	Bond	p-Isopropylphenyl	OMe	OMe	CH	O	O	
II-10	OH	Bond	Benzyl	OMe	OMe	CH	O	O	
II-11	OH	CH=CH	Ethyl	OMe	OMe	CH	O	O	
II-12	OH	CH=CH	(CH ₃) ₂ -CH ₂ -CH ₂	OMe	OMe	CH	O	O	
II-13	OH	CH=CH	Cyclopropylmethylene [sic]	OMe	OMe	CH	O	O	
II-14	OH	CH=CH	Methyl	OMe	O-CH ₂ -CH ₂ -C		O	O	
II-15	OH	CH ₂ -CH ₂	Ethyl	OMe	O-CH=CH-C		O	O	
II-16	OH	CH ₂ =CH ₂	Methyl	OMe	CH ₂ -CH ₂ -CH ₂ -C		O	O	
II-17	OH	Bond	Methyl	OMe	CH ₂ -CH ₂ -CH ₂ -C		O	O	147

Example 35

Receptor binding data were measured by the binding assay described above for the compounds listed below.

The results are shown in Table 2 [sic].

Table 2 [sic]

10

Receptor binding data (K_i values)

	Compound	ET_A [nM]	ET_B [nM]
15			
	I-2	6	34
	I-29	86	180
	I-5	12	160
20	I-4	7	2500
	I-87	1	57
	I-89	86	9300
	I-103	0.4	29
	I-107	3	485
25	I-12	19	1700
	I-26	23	2000
	I-23	209	1100
	I-47	150	1500
30	I-60	33	970
	I-96	0.6	56
	II-3	107	7300
35	II-1	28	2300

40

45